

Evidence for unusual events in high energy cosmic ray interactions

A.K. Argynova ^a, T.N. Kvochkina ^b, A.A. Loktionov ^{a 1} N.I. Kochelev^{c2}, A. V. Kropivnitskaya ^d, A.A Rostovtsev,^d

^a *Institute of Physics and Technology, Almaty, 480082, Kazakhstan*

^b *Institute of Nuclear Physics, Almaty, 480082, Kazakhstan*

^c *Joint Institute for Nuclear Research, Dubna, Moscow region, 141980 Russia*

^d *Institute for Theoretical and Experimental Physics, B.Chermushkinskaya 25, 117259, Moscow, Russia*

Abstract

Anomalous events with a large transverse momenta P_T were found for all secondary particles in an analysis of experimental data on cosmic ray interactions at $E_0 > 5 \cdot 10^{13} \text{ eV}$. In order to clarify the underlying mechanism of these events, the intermittency analysis of interactions with both large and usual transverse momentum was performed. It is shown that behavior of intermittency indices $\varphi(q)$ as a function of rank factorial momentum q for such interactions is very peculiar. The indices show slight growth for the events with large transverse momenta for all secondary particles in contrast with the fast growth for events with normal average transverse momenta for final particles. The significant difference in the behavior of intermittency indices gives strong evidence on the existence of the new class of hadron interactions with large P_T for all secondary particles in the event. A possible interpretation of these anomalous events as a manifestation of Color Glass Condensate in high energy hadron-hadron interactions is given.

¹loction@satsun.sci.kz

²kochelev@theor.jinr.ru

1 Introduction

Investigation of the properties of quark-gluon interactions at TEV region of energy is very important task, especially in the light of future experiments at LHC. One of the important goals of these experiments is clarification of the structure of high density/temperature quark-gluon matter which can be produced in heavy ion collisions. At present, cosmic ray experiments give the unique information on possible interesting effects in strong interaction of hadrons, which could be carefully investigated at LHC.

A possible manifestation of Color Glass Condensate (CGC) in high energy collisions is now widely under discussion (see recent reviews [1]). Phenomena of CGC arise in the high parton density regime of QCD where the typical virtualities of quarks and gluons are not very large and thus their interaction is not very strong. One of the most promising microscopical candidates for CGC is the sphaleron state [2] which is closely related to the instanton model of QCD vacuum (see reviews [3]) The feature of the sphaleron driven multiparticle production is rather high multiplicity in the events and large transverse momentum for all secondary particle.

In this Letter we will discuss a possible evidence for the formation of CGC in high energy ($E_0 > 5 \cdot 10^{13} eV$) cosmic ray interactions with nuclei in the stratosphere. An analysis of behavior of intermittency indices for unusual type of the events, where all the secondary particles were produced with a large transverse momenta $P_T \sim 1 GeV/c$ [4] is performed. An interesting feature of these events is a large difference in P_T distribution of the secondary particles compared to their distribution in the "usual" events, which have $P_T \approx 0.2 GeV/c$. Similar events were observed by JACCE [5] and Concorde [6] Collaborations in emulsion chambers. In spite of the fact that the first data on existence of "anomalous" events were obtained about twenty years ago, the underlying mechanism of hadron-hadron interaction, which is responsible for such events, is not understood so far. A promising way in investigation of dynamics of multiparticle production is the study of structure of the density fluctuations in the events through calculation of so-called intermittency indices [7]. The values of the indices are very sensitive to the microscopical mechanism of the hadron final state formation and, as we will show below, are very different for "anomalous" and "usual" cosmic rays events.

2 Experimental data

In this Section we present a new analysis of the experimental data, obtained in cosmic ray interactions at $E_0 > 5 \cdot 10^{13} eV$, which were observed in X -ray film - emulsion chambers exposed in the stratosphere at elevations $> 25 km$. Each chamber contained target and gamma blocks. The target blocks were gathered of many layers of heavy or light material interplead with thin nuclear emulsions; gamma blocks contained 4 - 10 lead plates, $5 mm$ thickness each with nuclear emulsions, and X -ray films between them. The details of the construction of the experimental chambers are described in [4].

For each event in the target block we determined the number of charged secondary particles (n_{ch}), the number of secondary γ - quanta (n_γ), the angles of emission of each secondary particle in the laboratory frame, Θ_{ch} , and, Θ_γ , an energy and a transverse momentum of each over threshold γ - quantum E_γ and $P_{T\gamma}$ (the threshold for the electromagnetic cascade in X -ray films was $2 TeV$ and in nuclear emulsion - $0.05 TeV$). We estimate the deviation of the accuracy for $P_{T\gamma}$ not greater than 30%. The number of secondary charged particles and their angular distribution provide an estimation of the energy of the primary particles which produce interactions in the target block of the chamber. The analysis of measured values of transverse momentum of each photon in events with $\sum E_\gamma > 10 TeV$ showed that 7 of them (index " $P_T = 1$ ") have rather different properties in compare with an others. Thus, the transverse momenta of most photons in these 7 events were several times larger than the average transverse momentum of secondary photons in the "usual" type events with $\langle P_{T\gamma} \rangle \sim 0.2 GeV/c$ (index " $P_T = 0$ ").

The integral distribution of transverse momentum of all the secondary photons is given in Fig.1 and can be described by the superposition of two exponents:

$$N_\gamma(> p_{t\gamma}) = A_1 \exp(-p_{t\gamma}/p_{01}) + A_2 \exp(-p_{t\gamma}/p_{02}), \quad (1)$$

where $p_{01} \sim 0.2 GeV/c$ for the "usual" events and $p_{02} \sim 0.8 GeV/c$ for the "anomalous" ones. The main feature of these 7 "anomalous" events is that most of the secondary photons have transverse momentum $p_{t\gamma} \geq 0.5 GeV/c$. The difference in characteristics between the two classes of events in Fig.1 is large large compared to the errors in the estimation of E_γ and can not be explained by loss of the underthreshold photons and statistical fluctuations.

3 Analysis of the pseudo-rapidity distributions of the secondary charged particles

At first glance, both types of events have a similar pattern. Thus, a comparison of χ^2/NDF fit with Gaussian distribution for events with higher multiplicity is given in table 1. The histograms of the pseudo-rapidity distributions of the secondary charged particles for these events are presented on Fig. 2. All events show approximately similar behavior.

Table 1. Multiplicities and χ^2/NDF of pseudo-rapidity distribution for large and small P_T events

N	small P_T events		large P_T events	
	NCH	χ^2	NCH	χ^2
1	290	1.99	675	3.09
2	239	2.19	432	1.32
3	175	1.24	235	2.42
4	174	3.16	182	2.25

4 Analysis of the structure of correlations

The analysis of the scaled factorial moments of the multiplicity distribution to isolate the dynamical density fluctuations from the background of statistical fluctuations was proposed in Ref. [8]:

$$\langle F_q(\delta) \rangle = \frac{\langle n^{[q]} \rangle}{\langle n \rangle^q} \quad (2)$$

where $\langle n^{[q]} \rangle = \langle n_m(n_m - 1) \dots (n_m - q + 1) \rangle$.

In the presence of self-similar fluctuations of different sizes, the dependence of the moment $\langle F_q \rangle$ on the size of the phase-space bin follows the power law:

$$\langle F_q \rangle = (1/\delta)^{\varphi_q} \quad (3)$$

for $\delta \rightarrow 0$. The positive constant φ_q is the so-called intermittency exponent.

The distribution (2) is not a smooth function, i.e. it contains sharp spikes and holes between particles in the phase-space. The observation of

the power law in a sufficiently large range of scales δ indicates a self-similar fractal structure of the short range particle density fluctuations.

In the opposite case of a smooth distribution (probability density is continuous), the factorial moments are

$$\langle F_q \rangle \sim Const. \quad (4)$$

It should be mentioned that the high order moments resolve the tail of the multiplicity distribution. Thus they are very sensitive to the density fluctuations at various scales of δ used in the analysis. Therefore, the study of fluctuations at various values of δ , especially at small δ and large q can improve our phenomenological understanding of multiparticle production processes.

The particle multiplicity distribution is usually studied for a sequence of phase space domains δ by consecutive subdivision of an initial region Δ in M equal subdomains:

$$\delta = \Delta/M$$

In every subdomain m ($m = 1, \dots, M$) n_m is the multiplicity of that bin.

In order to improve the statistical accuracy in the experimental estimation of factorial moments, F_q 's of individual cells (1) are averaged over events and over M cells ("vertical analysis"). The moments averaged vertically (*i.e.* over events) can be defined as a double average

$$\langle F_q \rangle = \frac{1}{M} \sum_m \frac{1}{N_{evt}} \sum_{evt} \frac{\langle n_m^{[q]} \rangle}{\langle n_m \rangle^q}, \quad (5)$$

where $\langle n_m \rangle = \frac{1}{N_{evt}} \sum_{evt} n_m$ is the multiplicity in bin m .

We use a modified method of vertically averaging [12] in which moments are averaged at the starting point of the original region Δ location

$$\langle F_q \rangle = \frac{1}{M} \sum_m \frac{1}{N_{step}} \sum_{step} \frac{1}{N_{evt}} \sum_{evt} \frac{\langle n_m^{[q]} \rangle}{\langle n_m \rangle^q}, \quad (6)$$

where N_{step} is the number of small ($step/\Delta \ll 1$) steps of the starting position of the original region Δ in the area of pionization. The pseudo-rapidity $\eta = -\ln tg\Theta/2$ of the secondary particles charge is used as the basic variable. The initial value of Δ is 4.0 and $M = 40$. More detailed analysis has been performed with $\Delta = 3.5, 3.8$, and $M = 35, 38$, respectively.

5 The results

The results are presented in Table 2 and in Fig.3-5. The values of $\ln F_q$'s as a function of $-\ln/\delta\eta$ for order $q = 2$ to $q = 8$, are plotted for small p_t events in Fig.3 and for events with large P_T in Fig.4.

For small p_t events in Fig.3 the rise of $\ln F_q$ when increasing $-\ln \delta\eta$ takes place for all q . For large P_T events (Fig.4) at all q any significant $\delta\eta$ dependence is absent.

The slopes φ_q in the region $0.1 \leq \ln \delta\eta \leq 1.0$ indicated by the solid lines in the figures are given in Table 2. Values of φ_q are essentially larger for the events with small P_T compared to the large P_T events.

The comparison of the slopes φ_q for both types of events is shown in Fig.5. For small P_T events the data are consistent with the intermittent behaviour (2), while for "anomalous" events the values of the slopes and their changing as function of q are essentially smaller.

Table 2. Slopes φ_q fitted over $0.1 \leq \delta\eta \leq 1.0$ for large and small P_T events

q	small P_T events	large P_T events
	φ_q	φ_q
2	0.100 ± 0.004	0.068 ± 0.005
3	0.260 ± 0.014	0.095 ± 0.010
4	0.310 ± 0.027	0.094 ± 0.016
5	0.51 ± 0.05	0.08 ± 0.02
6	0.66 ± 0.06	0.10 ± 0.03
7	0.77 ± 0.09	0.11 ± 0.04
8	1.29 ± 0.11	0.13 ± 0.06

6 Possible explanation of anomalous events

Self-similar particle-density fluctuations have been observed in various reactions [7]. Thus, the approximate scaling in respect to decreasing of domain size was found. In spite of the strong experimental evidence, the effect is still rather far from understanding based on fundamental QCD theory. One of the possible contributions, which might give strong influence on behavior of

scaling moments, is coming from Bose-Einstein correlations. However, these correlation should not be related to some conventional static source [9].

There is a clear indication on P_T - dependence of intermittency in different processes [10], [11]. An analysis for particles with P_T smaller or larger than $0.15 \div 0.3 GeV/c$ in the same events reveals a strong sensitivity to the value of transverse momentum. The results show that the slopes φ_q increase by a factor $2 \div 4$ for value of cut $P_T = 0.15 GeV$. A similar but not that strong effect is observed if the P_T cut is moved to $0.30 GeV/c$.

Our results for small P_T events confirm the power-law behavior given by Eq.2. On a contrary, for the large P_T events, expression (3) looks as a very promising candidate for the behavior of the intermittency exponents.

The results on P_T distribution, presented in Fig. 1, and the results related to the intermittency (Fig. 3 - 5 and Tables 1, 2), strongly indicate the existence of a new class of events with large P_T for all secondary particles. We should mention that intermittency dependence for the "usual" events might be understood as a result of perturbative QCD evolution of partonic cascade at high energy. In this case the independent hadronization of the quarks and gluons in partonic chain should lead to a significant intermittency patterns [7]. For the "anomalous" large P_T events, we found that the behavior of the factorial moments is very different compared to that of the "usual" events. From our point of view this is very important result. It seems that in this case the perturbative QCD can not describe the data and a new mechanism for secondary particle production should be involved. One of the possible candidates for this mechanism is the contribution from nonperturbative partonic interactions at high energies, related to the Color Glass Condensate formation at large parton density [1]. In this case the process of particle production in very high energy nucleus-nucleus interactions induced by cosmic rays looks like simultaneous burst of all interacted regions in the overlapped cosmic and target nuclei. If the CGC is identified with the QCD sphaleron state [2], which describes the particle production induced by strong non-perturbative fluctuations of vacuum gluon fields, instantons [3], than the transverse momentum of produced particles should be roughly $P_T(large) \approx 1/\rho_c \approx 0.6 GeV$, where $\rho_c \approx 0.3 fm$ is the average size of instantons in QCD vacuum. This value of transverse momentum is much larger than the value of transverse momentum $P_T(small) \approx 1/R_{conf} \approx 0.2 GeV$ ($R_{conf} \approx 1 fm$ is the confinement size) of secondary particles in "usual" events.

7 Conclusion

In summary, a strong indication of the existence of a new class of hadron-hadron interactions at high energies is found. A possible explanation of these interactions is a manifestation of the Color Glass Condensate formation in the cosmic ray induced events. It would be interesting to search for events with similar properties at current (HERA, RHIC, TEVATRON) and future (LHC, TESLA) colliders.

Acknowledgements

We are grateful to A.E.Dorokhov for useful discussion. This work was partially supported by grants RFBR-03-02-17291 and RFBR-04-02-16445.

References

- [1] R.Venugopalan, hep-ph/0412396; E. Iancu and R. Venugopalan, hep-ph/0303204; E. Iancu, A. Leonidov and L. McLerran, hep-ph/0202270.
- [2] F. Schrempp and A. Utermann, hep-ph/0407146; hep-ph/0401137; Phys. Lett. **543** (2002) 197.
- [3] T. Schäfer and E.V. Shuryak, Rev.Mod.Phys. **70** (1998) 1323; A.E.Dorokhov, N.I.Kochelev and Yu.A. Zubov, Int.Jour. of Mod. Phys. **A8** (1993) 603; D. Diakonov, Prog. Par. Nucl. Phys. **51** (2003) 173.
- [4] N. Dobrotin, V. Ivanenko, T. Kvochkina. Proc.16-th ICRC, Kyoto,**v.7** (1979) 268.; A. Apanasenko, N. Dobrotin, V. Ivanenko, T. Kvochkina et al. Proc. 21-th ICRC, Adelaide, **v.8** (1990) 112; 141.
- [5] JACCE Coll. T. Burnett et al, Phys.Rev.Lett., **bf 57** (1986) 3249; W.V.Jones et al. Ann. Rev. Nucl. Part. Sci., **37** (1987) 71.
- [6] Concorde Coll. J.Phys.G. **12** (1986) 12; J.Phys.G. **14** (1988) 503.
- [7] E.A. De Wolf, I.M. Dremin, W. Kittel, Phys. Rept. **270** (1996) 1; I.M. Dremin and J.W. Gary, Phys. Rept. **349** (2001) 301; I.M. Dremin, Usp. Fiz. Nauk. **172** (2002) 551, hep-ph/0203024.

- [8] A.Bialas and R.Peschanski, Nucl. Phys. **B273** (1986) 703; Nucl. Phys. **B308** (1988) 857.
- [9] A. Bialas, Acta Phys. Pol. **B23** (1992) 561.
- [10] N.N. Biswas et al., Phys. Rev. Lett. **37** (1976) 175.
- [11] I.V. Ajinenko et al. (NA 22), Z. Phys. **C58** (1993) 357.
- [12] A.Kh. Argynova et al., Report at 22-nd Intern Nathiagali Summer College on Physics and Contemporary Needs, Islamabad, Pakistan, 28 July -9 August, 1997; Report at 25th Cosmic Ray Conf., Moscow, 22-27 June, 1998.

FIGURE CAPTIONS

Fig. 1. P_T distribution of secondary particles in two types of events.

Fig. 2. Pseudo-rapidity distribution of events with small and large P_T .

Fig. 3. $\ln \langle F_q \rangle$ as a function of $-\ln \delta \eta$ for the group of small P_T events.

Fig. 4. $\ln \langle F_q \rangle$ as a function of $-\ln \delta$ for the group of large P_T events.

Fig. 5. The comparison of the slopes φ_q as function of the moment order q for two types of events.






Fig. 1.

$P_T 0$

$P_T 1$



174

182





175

235



239

432



290

675





Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.